

# RESTRICTED 內部文件

香港考試局

HONG KONG EXAMINATIONS AUTHORITY

一九八七年香港中學會考

HONG KONG CERTIFICATE OF EDUCATION EXAMINATION, 1987

附加數學 (卷二)

ADDITIONAL MATHEMATICS (Paper II)

評卷參考

MARKING SCHEME

這份內部文件，只限閱卷員參閱，不得以任何形式翻印。

This is a restricted document.

It is meant for use by markers of this paper for marking purposes only.

Reproduction in any form is strictly prohibited.

請在學校任教之閱卷員特別留意

本評卷參考並非標準答案，故極不宜落於學生手中，以免引起誤會。

遇有學生求取此文件時，閱卷員應嚴予拒絕。閱卷員在任何情況下披露本評卷參考內容，均有違閱卷員守則及「一九七七年香港考試局法例」。

Special Note for Teacher Markers

It is highly undesirable that this marking scheme should fall into the hands of students. They are likely to regard it as a set of model answers, which it certainly is not.

Markers should therefore resist pleas from their students to have access to this document. Making it available would constitute misconduct on the part of the marker and is, moreover, in breach of the 1977 Hong Kong Examinations Authority Ordinance.

© 香港考試局 保留版權  
Hong Kong Examinations Authority  
All Rights Reserved 1987

RESTRICTED 內部文件


Provided by dse.life

SOLUTIONS	MARKS	REMARKS
<p>1. <math>(1 + x + x^2)^n</math>  <math>= [1 + x(1 + x)]^n</math> .....  <math>= 1 + nx(1 + x) + \frac{n(n-1)}{2} (x^2)(1 + x)^2 + \dots</math></p> <p>Coeff. of <math>x^2 = n + \frac{n(n-1)}{2}</math>  <math>= 21</math></p> <p><math>n^2 + n - 42 = 0</math>  <math>(n - 6)(n + 7) = 0</math> .....  <math>n = 6</math> or <math>-7</math> (rejected) .. .. .</p>	<p>1M  2A  1A 1A <hr/>5</p>	
<p>2. For <math>n = 1</math>, L.H.S. = <math>1/4</math>  R.H.S. = <math>1/4 =</math> L.H.S. ....</p> <p>Assume equality holds for some integer <math>k</math>.</p> <p>For <math>n = k + 1</math>,</p> <p>L.H.S. = <math>\frac{1}{(1)(4)} + \frac{1}{(4)(7)} + \dots + \frac{1}{(3k-2)(3k+1)} + \frac{1}{(3k+1)(3k+4)}</math>  <math>= \frac{k}{3k+1} + \frac{1}{(3k+1)(3k+4)}</math>  <math>= \frac{3k^2 + 4k + 1}{(3k+1)(3k+4)}</math>  <math>= \frac{k+1}{3k+4}</math> .....  <math>=</math> R.H.S.</p> <p>Therefore equality holds also for <math>n = k + 1</math>.  mathematical induction, equality holds for all  positive integers <math>n</math>.</p>	<p>1  1  1  1  <hr/>5</p>	<p>Award this mark only if the candidate has scored the first four marks.</p>
<p>3. Let the slope of the required line be <math>m</math>.</p> <p><math>\frac{m-3}{1+(m)(3)} = \pm \frac{1}{2}</math></p> <p><math>2(m-3) = 3m+1</math> or <math>2(m-3) = -(3m+1)</math></p> <p><math>m = -7</math> or <math>m = 1</math></p> <p><math>\frac{y-2}{x-1} = -7</math>      <math>\frac{y-2}{x-1} = 1</math></p> <p><math>7x + y - 9 = 0</math>      <math>x - y + 1 = 0</math></p>	<p>1A+1  1A+1A  <hr/>1A 5</p>	<p>1A for formula (excl. <math>\pm</math>) 1 for <math>\pm</math></p> <p>For both equations</p>

SOLUTIONS	MARKS	REMARKS
<p>4. Put <math>x = \sin\theta</math>  <math>dx = \cos\theta d\theta</math> .....</p> <p><math>x = 0, \theta = 0</math> )  <math>x = \frac{1}{2}, \theta = \frac{\pi}{6}</math> ) .....</p> $\int_0^{\frac{1}{2}} \frac{2x^2}{\sqrt{1-x^2}} dx = \int_0^{\frac{\pi}{6}} \frac{2\sin^2\theta}{\sqrt{1-\sin^2\theta}} \cos\theta d\theta$ $= \int_0^{\frac{\pi}{6}} 2 \sin^2\theta d\theta$ $= \int_0^{\frac{\pi}{6}} (1 - \cos 2\theta) d\theta$ $= \left[ \theta - \frac{1}{2} \sin 2\theta \right]_0^{\frac{\pi}{6}}$ $= \frac{\pi}{6} - \frac{\sqrt{3}}{4} \quad (0.0906)$	<p>1A</p> <p>1A</p> <p>1A</p> <p>1A</p> <p>1A</p> <p><u>1A</u></p> <p><u>6</u></p>	<p>)</p> <p>) for integrand</p> <p>)</p>
<p>5. <math>y = \int (3x^2 - 2)(x^3 - 2x + 1)^{\frac{1}{3}} dx</math></p> <p>put <math>u = x^3 - 2x + 1</math>  <math>du = (3x^2 - 2) dx</math> .....</p> $y = \int u^{\frac{1}{3}} du$ $y = \frac{3}{4} u^{\frac{4}{3}} + c$ $y = \frac{3}{4} (x^3 - 2x + 1)^{\frac{4}{3}} + c$ <p>sub. <math>x = 0, y = 0</math></p> $c = -\frac{3}{4}$ $y = \frac{3}{4} (x^3 - 2x + 1)^{\frac{4}{3}} - \frac{3}{4}$	<p>1M</p> <p>1A</p> <p>1A</p> <p>1A</p> <p>1M</p> <p><u>1A</u></p> <p><u>6</u></p>	<p>Do not award this mark if c is missing.</p>
<p>6. <math>\sin 3\theta = \sin 2\theta \cos \theta + \cos 2\theta \sin \theta</math>  <math>= 2\sin \theta \cos^2 \theta + (1 - 2\sin^2 \theta) \sin \theta</math>  <math>= 3\sin \theta - 4\sin^3 \theta</math></p> <p>Put <math>x = \sin \theta</math> .....</p> $8x^3 - 6x + 1 = 0$ $8\sin^3 \theta - 6\sin \theta + 1 = 0$ $2(4\sin^3 \theta - 3\sin \theta) + 1 = 0$ $2\sin 3\theta = 1$ $\sin 3\theta = \frac{1}{2}$ $3\theta = 180n^\circ + (-1)^n 30^\circ$ $\theta = 60n^\circ + (-1)^n 10^\circ$ $= 10^\circ, 50^\circ, 130^\circ, 170^\circ, 250^\circ, \dots$ <p><math>x = \sin 10^\circ, \sin 50^\circ, \sin 250^\circ</math>  <math>= 0.17, 0.77, -0.94</math></p>	<p>1A</p> <p>1A</p> <p>1M</p> <p>1A</p> <p>1A+1A</p> <p><u>6</u></p>	<p><math>(\cos \theta + i \sin \theta)^3</math>  <math>= \cos 3\theta + i \sin 3\theta</math> 1A</p> <p>⋮</p> <p><math>\sin 3\theta = 3\sin \theta - 4\sin^3 \theta</math> 1A</p> <p>2 correct answers 1A          3 correct answers 2A</p>

SOLUTIONS	MARKS	REMARKS
7. Tangents are of the form $y = 2x + k$	1A	<u>Alternative Solution:</u>
Sub. in $x^2 - y^2 = 3$ .....	1M	Diff. $x^2 - y^2 = 3$ <span style="float: right;">1M</span>
$x^2 - (2x + k)^2 = 3$		$2x - 2yy' = 0$ <span style="float: right;">1A</span>
$-3x^2 - 4kx - k^2 = 3$		$y' = \frac{x}{y}$
$3x^2 + 4kx + k^2 + 3 = 0$	1A	$\frac{x}{y} = 2$ <span style="float: right;">1A</span>
For tangents, $\Delta = 0$		$x = 2y$
$16k^2 - 4(3)(k^2 + 3) = 0$ .....	1M	Sub. in $x^2 - y^2 = 3$ <span style="float: right;">1M</span>
$k^2 = 9$		$3y^2 = 3$
$k = \pm 3$	1A+1A	$y = \pm 1$
Equations of tangents $y = 2x + 3$ and $y = 2x - 3$	<u>6</u>	$x = \pm 2$
		$y = 2x-3$ and $y = 2x+3$ <span style="float: right;">1A+1A</span>
<p><u>Alternative Solution:</u></p> <p>Eq. of tangent: <math>x_1x - y_1y = 3</math> .....</p> <p style="padding-left: 40px;">slope = <math>\frac{x_1}{y_1}</math> <span style="float: right;">1A</span></p> <p style="padding-left: 40px;"><math>\frac{x_1}{y_1} = 2</math> .....</p> <p style="padding-left: 40px;"><math>\frac{x_1}{y_1} = 2</math> <span style="float: right;">1A</span></p> <p>etc.</p>		


SOLUTIONS	MARKS	REMARKS
8. (a) $du = \sec^2 x dx$ ,	1A	
$\int \tan^{n-2} x \sec^2 x dx = \int u^{n-2} du \dots\dots\dots$	1A	
$= \frac{\tan^{n-1} x}{n-1} + c$	<u>1A+1A</u> 4	1A for c
(b) (i) $\int_0^{\frac{\pi}{4}} \tan^n x dx = \int_0^{\frac{\pi}{4}} \tan^{n-2} x \tan^2 x dx$		
$= \int_0^{\frac{\pi}{4}} \tan^{n-2} x (\sec^2 x - 1) dx$	1A	
$= \int_0^{\frac{\pi}{4}} \tan^{n-2} x \sec^2 x dx - \int_0^{\frac{\pi}{4}} \tan^{n-2} x dx$	1M	
$= \left[ \frac{\tan^{n-1} x}{(n-1)} \right]_0^{\frac{\pi}{4}} - \int_0^{\frac{\pi}{4}} \tan^{n-2} x dx$	1M	1M for using (a)
$= \frac{1}{n-1} - \int_0^{\frac{\pi}{4}} \tan^{n-2} x dx$	1	<u>Alternative Solution:</u>
(ii) $I_0 = \int_0^{\frac{\pi}{4}} dx = \frac{\pi}{4}$ or $I_2 = 1 - \frac{\pi}{4} \dots\dots\dots$	1A	$\int_0^{\frac{\pi}{4}} \tan^6 x dx$
$I_6 = \int_0^{\frac{\pi}{4}} \tan^6 x dx = \left( \frac{1}{5} - I_4 \right)$	2A	$= \int_0^{\frac{\pi}{4}} \tan^4 x (\sec^2 x - 1) dx$ 1A
$I_4 = \left( \frac{1}{3} - I_2 \right)$	1A	⋮
$I_6 = \left[ \frac{1}{5} - \frac{1}{3} + 1 - I_0 \right]$		$= \left[ \frac{1}{5} \tan^5 x - \frac{1}{3} \tan^3 x + (\tan x - x) \right]_0^{\frac{\pi}{4}}$
$= \left( \frac{13}{15} - \frac{\pi}{4} \right)$ or 0.0813	1A	1A+1A+1A
	<u>9</u>	$= \frac{13}{15} - \frac{\pi}{4} \dots\dots\dots$ 1A
(c) Putting $x = -v$ $\dots\dots\dots$	1A	<u>Alternative Solution:</u>
$dx = -dv$	1A	$\int_{-\frac{\pi}{4}}^0 \tan^6 x dx$
$x = 0, v = 0$ )		0
$x = -\frac{\pi}{4}, v = \frac{\pi}{4}$ ) $\dots\dots\dots$	1A	$= \left[ \frac{1}{5} \tan^5 x - \frac{1}{3} \tan^3 x + (\tan x - x) \right]_{-\frac{\pi}{4}}^0$
$\int_{-\frac{\pi}{4}}^0 \tan^6 x dx = \int_{\frac{\pi}{4}}^0 \tan^6(-v) (-dv)$		⋮
$= \int_{\frac{\pi}{4}}^0 \tan^6 v dv$	1	$= \frac{13}{15} - \frac{\pi}{4} \dots\dots\dots$ 1A
$\int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \tan^6 x dx = \int_{-\frac{\pi}{4}}^0 \tan^6 x dx + \int_0^{\frac{\pi}{4}} \tan^6 x dx$	1A	$= \int_0^{\frac{\pi}{4}} \tan^6 x dx$ 1
$= 2 \int_0^{\frac{\pi}{4}} \tan^6 x dx \dots\dots\dots$	1A	
$= 2 \left( \frac{13}{15} - \frac{\pi}{4} \right)$ or 0.163	<u>1A</u> 7	

SOLUTIONS	MARKS	REMARKS
<p>9. (a)</p>  <p>Area of region I = <math>\int_0^s x^2 dx</math></p> $= \left[ \frac{x^3}{3} \right]_0^s$ $= \frac{s^3}{3}$ <p>Area of (shaded region + I + II)</p> $= \frac{1}{2}(s+t)(s^2+t^2)$ <p>Area of region II = <math>\frac{t^3}{3}</math> .....</p> <p>Shaded area = <math>\frac{1}{2}(s+t)(s^2+t^2) - \frac{1}{3}s^3 - \frac{1}{3}t^3</math></p> $= \frac{1}{6}(s^3 + 3s^2t + 3st^2 + t^3)$ $= \frac{1}{6}(s+t)^3$	<p>1A</p> <p>1A</p> <p>1A</p> <p>1A</p> <p>1M+1A</p> <p>1</p> <hr/> <p>7</p>	<p>Alternative Solution:</p> <p>ST : <math>\frac{y-s^2}{x-s} = \frac{s^2-t^2}{s-(-t)}</math></p> <p><math>y = (s-t)x + st</math> 1A</p> <p>Shaded area</p> $= \int_{-t}^s [(s-t)x + st - x^2] dx$ 1M+1A $= \left[ \frac{(s-t)x^2}{2} + stx - \frac{x^3}{3} \right]_{-t}^s$ 1A $= \frac{1}{6}(s^3 + 3s^2t + 3st^2 + t^3)$ $= \frac{1}{6}(s+t)^3$
<p>(b) (i) S, H, T are collinear.</p> $\frac{s^2-1}{s-0} = \frac{t^2-1}{-t-0}$ ..... $-s^2t + t = st^2 - s$ $s + t = st(t+s)$ $st = 1$ $t = \frac{1}{s}$ .....	<p>1M</p> <p>1</p>	<p>Sub. (0, 1) in eqt. of ST 1M</p> <p><math>1 = st</math></p> <p><math>t = \frac{1}{s}</math> .....</p>
<p>(ii) Shaded area <math>A = \frac{1}{6}(s + \frac{1}{s})^3</math></p> $\frac{dA}{ds} = \frac{1}{6}(3)(s + \frac{1}{s})^2(1 - \frac{1}{s^2})$ ..... $= 0$ <p><math>s = 1</math> or <math>-1</math> (rejected)</p> <p><math>\therefore s = 1</math> .....</p> <p><math>s &lt; 1, \frac{dA}{ds} &lt; 0</math> )</p> <p><math>s &gt; 1, \frac{dA}{ds} &gt; 0</math> ) .....</p> <p><math>\therefore s = 1</math> corresponds to a minimum A .</p>	<p>1A</p> <p>1A</p> <p>1M</p> <p>1A</p> <p>1M</p> <hr/> <p>7</p>	$\frac{d^2A}{ds^2} = \frac{1}{2} 2(s + \frac{1}{s})(1 - \frac{1}{s^2})^2$ $+ \frac{1}{2} (s + \frac{1}{s})^2 (\frac{2}{s^3})$ <p>When <math>s = 1, \frac{d^2A}{ds^2} &gt; 0</math> 1M</p>

SOLUTIONS	MARKS	REMARKS
<p>9. (c) For <math>s = 1</math>, ST is horizontal.</p> <p>Volume generated by region I = <math>\int_0^1 \pi y^2 dx</math></p> $= \pi \int_0^1 x^4 dx$ $= \pi \left[ \frac{x^5}{5} \right]_0^1$ $= \frac{1}{5} \pi \dots\dots\dots$ <p>Volume of cylinder = <math>\pi(1)^2(2)</math></p> <p>Required volume = <math>\pi(1)^2(2) - \frac{1}{5}\pi - \frac{1}{5}\pi</math></p> $= \frac{8\pi}{5} \text{ (or 5.03) } \dots\dots\dots$	<p>1M</p> <p>1A</p> <p>1A</p> <p>1M</p> <p><u>1A</u></p> <p><u>6</u></p>	<p>For <math>\int_a^b \pi y^2 dx</math></p>
<p><u>Alt. Solution</u></p> <p>Volume generated = <math>\int_a^b \pi(y_1^2 - y_2^2) dx</math></p> $= 2 \int_0^1 \pi(1 - x^4) dx$ $= 2\pi \left[ x - \frac{x^5}{5} \right]_0^1$ $= \frac{8\pi}{5} \text{ (or 5.03) } \dots\dots\dots$	<p>1M+1M</p> <p>2A</p> <p>1A</p> <p>1A</p>	<p>1M for <math>\int_a^b \pi y^2 dx</math></p>

SOLUTIONS	MARKS	REMARKS
<p>10.(a) PS = PN</p> $\sqrt{(x-1)^2 + y^2} = x+1$ $(x-1)^2 + y^2 = (x+1)^2$ $y^2 = 4x \dots\dots\dots$	<p>1M+1A +1A</p> <hr style="width: 50%; margin: 0 auto;"/> <p>1</p> <hr style="width: 50%; margin: 0 auto;"/> <p>4</p>	<p>1A for L.S. 1A for R.S.</p>
<p>(b) (i) <math>y = 2t</math> <math>x = t^2 \dots\dots\dots</math></p> <p>(ii)(1) PN // x-axis and PR bisects <math>\angle</math> SPN. <math>\therefore \angle</math> PRS = <math>\angle</math> RPS SR = SP = PN = <math>t^2 + 1 \dots\dots\dots</math> <math>\therefore</math> OR = SR - SO = <math>t^2 \dots\dots\dots</math> R is the point <math>(-t^2, 0)</math> <math>\therefore</math> the equation of PR is <math>y = \frac{2t-0}{t^2 - (-t^2)}(x+t^2) \dots\dots\dots</math> i.e. <math>x - ty + t^2 = 0</math></p>	<p>1A</p> <p>2A</p> <p>1A</p> <p>2A</p> <p>1M</p> <p>1</p>	<p><u>Alternative Solution:</u> PR intersects SN at M M is the mid-point of SN 3A <math>\therefore</math> M is the point <math>(0, t)</math> 2A PR : <math>\frac{y-t}{x-0} = \frac{2t-t}{t^2-0}</math> 1M <math>x - ty + t^2 = 0 \dots\dots\dots</math> 1</p>
<p><u>Alternative Solution:</u> PS : <math>\frac{y-0}{x-1} = \frac{2t}{t^2-1}</math> <math>2tx + (1-t^2)y - 2t = 0 \dots\dots\dots</math> PN : <math>y = 2t</math> PR is the angle bisector. Its equation is <math>\frac{y-2t}{\sqrt{1^2+0^2}} = \frac{2tx+(1-t^2)y-2t}{\sqrt{(2t)^2+(1-t^2)^2}}</math> <math>y-2t = \frac{2tx+(1-t^2)y-2t}{1+t^2} \dots\dots\dots</math> <math>x - ty + t^2 = 0 \dots\dots\dots</math></p>	<p>1M</p> <p>1A</p> <p>1A</p> <p>2M+1A</p> <p>1</p>	
<p>(2) Sub. <math>x = ty - t^2</math> in <math>y^2 = 4x \dots\dots\dots</math> <math>4(ty - t^2) = y^2</math> <math>y^2 - 4ty + 4t^2 = 0 \dots\dots\dots</math> <math>\Delta = (-4t)^2 - 4(4t^2) \quad 1M</math> <math>= 0 \quad 1A</math> <span style="border: 1px solid black; padding: 2px;"><math>(y-2t)^2 = 0</math></span> 2A <math>\therefore</math> it touches <math>y^2 = 4x</math> at P.</p>	<p>1M</p> <p>1A</p> <p>2A</p>	<p><u>Alternative Solution:</u> Differentiating <math>y^2 = 4x</math> 1M <math>y' = \frac{2}{y}</math> slope of tangent at P = <math>\frac{1}{t}</math> 1A Eqt. of tangent at P: <math>y - 2t = \frac{1}{t}(x - t^2) \dots\dots\dots</math> 1M <math>x - ty + t^2 = 0 \dots\dots\dots</math> 1A which is the eqt. of PR. <math>\therefore</math> PR touches <math>y^2 = 4x</math> at P</p>
<p>(3) R is the point <math>(-t^2, 0)</math> _____ P is the point <math>(t^2, 2t)</math> Mid-point of PR is <math>(0, t) \dots\dots\dots</math> Equation of locus is <math>x = 0</math>.</p>	<p>1A</p> <p>1A</p> <p>2A</p> <hr style="width: 50%; margin: 0 auto;"/> <p>16</p>	



SOLUTIONS	MARKS	REMARKS
<p>11.(a) (i) <math>x^2 + y^2 - 16x - 4y + 64 = 0</math></p> <p>Put <math>y = 0</math>,</p> $x^2 - 16x + 64 = 0$ $(x - 8)^2 = 0 \text{ or } \Delta = (-16)^2 - 4(64) = 0$ $x = 8$ <p>Therefore <math>C_1</math> touches the x-axis at A</p>	<p>1M</p> <p>1A</p>	<p>Centre = (8, 2) )</p> <p>radius = 2 )</p> <p>Distance from centre to x-axis = radius 1</p> <p><math>C_1</math> touches the x-axis at A</p>
<p>(ii) Let equation of OH be <math>y = mx</math></p> <p>Sub. in equation of <math>C_1</math></p> $x^2 + m^2x^2 - 16x - 4mx + 64 = 0$ $(1 + m^2)x^2 - 4(m + 4)x + 64 = 0$ <p>For tangents,</p> $16(m + 4)^2 - (4)(64)(1 + m^2) = 0$ $m^2 + 8m + 16 - 16m^2 - 16 = 0$ $15m^2 - 8m = 0$ <p><math>m = 0</math> or <math>\frac{8}{15}</math></p> <p>OH : <math>y = \frac{8}{15}x</math> .....</p>	<p>1A</p> <p>1A</p> <p>1M+1.</p> <p>1M</p> <p>1A</p> <p>1A</p>	<p><u>Alternative Solution:</u></p> <p>OH: <math>y = mx</math> 1.</p> <p><math>C_1</math>: centre = (8,2) radius = 2</p> <p><math>\frac{8m - 2}{\sqrt{1 + m^2}} = \pm 2</math> (<math>\pm</math> optional) 1M+1.</p> <p><math>(4m - 1)^2 = 1 + m^2</math></p> <p><math>15m^2 - 8m = 0</math> 1.</p> <p><math>m = 0</math> or <math>\frac{8}{15}</math></p> <p>OH: <math>y = \frac{8}{15}x</math> 1.</p>
<p><u>Alternative Solution:</u></p> <div style="display: flex; align-items: center;">  <div style="margin-left: 20px;"> <p>Eqn. of OH: <math>y = mx</math> 1A</p> <p><math>\tan\theta = \frac{2}{8} = \frac{1}{4}</math> 1A</p> <p><math>m = \tan \angle AOH</math></p> <p><math>= \tan 2\theta</math> 1A</p> <p><math>= \frac{2\tan\theta}{1 - \tan^2\theta}</math> 1M</p> <p><math>= \frac{8}{15}</math> ..... 1A</p> </div> </div>		
<p>(iii) Let coordinates of H be <math>(8, y_1)</math></p> <p>Sub. in equation of OH</p> $y_1 = \frac{64}{15}$ <p>Equation of BH : <math>\frac{y - 0}{x - 16} = \frac{\frac{64}{15} - 0}{8 - 16}</math></p> $\frac{y}{x - 16} = -\frac{8}{15}$ $y = -\frac{8}{15}x + \frac{128}{15}$ $8x + 15y - 128 = 0$	<p>1A</p> <p>1M</p> <p>1A</p> <p>1M</p> <p>1A</p>	<p><u>Alternative Solution:</u></p> <p>By symmetry or <math>\angle HOB = \angle OBH</math>.</p> <p>Slope of BH = <math>\tan(180^\circ - \angle BOH)</math></p> <p><math>= -\tan \angle BOH</math></p> <p><math>= -\frac{8}{15}</math> 3A</p> <p>BH: <math>\frac{y - 0}{x - 16} = -\frac{8}{15}</math> 1M</p> <p><math>8x + 15y - 128 = 0</math> 1A</p>
	<p><u>12</u></p>	

SOLUTIONS	MARKS	REMARKS
11.(b) (i) Sub. (8, 0) in equation of $C_2$ $64 - 128 + c = 0$ $c = 64$ $C_2$ touches $4x + 3y = 0$ Sub. in $C_2$ $x^2 + \frac{16}{9}x^2 - 16x - \frac{8f}{3}x + 64 = 0$ $25x^2 - (144 + 24f)x + (9)(64) = 0$ For tangents, $(144 + 24f)^2 - 4(25)(9)(64) = 0$ $f = 4 \text{ or } -16 \dots\dots\dots$ Rejecting $f = -16$ , $f = 4$	1M  1A           1M 1A    1A   1A   1A   1M   2A  8	Put $y = 0$ in eqt. of $C_2$ $x^2 - 16x + c = 0$ $\Delta = 16^2 - 4c = 0 \dots\dots\dots 1A$ $c = 64 \dots\dots\dots 1A$ <u>Alternative Solution:</u> OK is tangent. Centre of $C_2 = (8, -f)$ radius = $f$ $\therefore \frac{4(8) - 3(f)}{\sqrt{4^2 + 3^2}} = \pm f \dots\dots\dots 1M$ $32 - 3f = \pm 5f$ $f = 4 \text{ or } -16 \dots\dots\dots 1A$ Rejecting $f = -16$ $f = 4 \dots\dots\dots 1A$ <u>Alt. Solution:</u> $K = (8, k)$ Sub. in $4x+3y = 0$ <span style="float: right;">1M</span> $k = -\frac{32}{3}$ $\frac{\Delta_{OBH}}{\Delta_{OBK}} = \frac{AH}{AK}$ $= \frac{64/15}{32/3}$ $= \frac{2}{5} \dots\dots\dots 2A$
(ii) $\frac{\Delta_{OBH}}{\Delta_{OBK}} = \frac{\frac{1}{2}(OB)(AH)}{\frac{1}{2}(OB)(AK)}$ $= \frac{AH}{AK}$ $= \frac{AH/OA}{AK/OA} \dots\dots\dots$ $= \frac{8/15}{4/3}$ $= \frac{2}{5}$	1M           2A  8	
<u>Alternative Solution:</u> $\Delta_{OBH} = \frac{1}{2} (16) \left( \frac{64}{15} \right) \dots\dots\dots 1A$ $\Delta_{OBK} = \frac{1}{2} (16) \left( \frac{32}{3} \right) \dots\dots\dots 1A$ $\frac{\Delta_{OBH}}{\Delta_{OBK}} = \frac{2}{5} \dots\dots\dots 1A$		

SOLUTIONS		MARKS	REMARKS
12.(a)(i)	$7\sin\theta - 24\cos\theta$ $= \sqrt{7^2+24^2} \left( \frac{7}{\sqrt{7^2+24^2}} \sin\theta - \frac{24}{\sqrt{7^2+24^2}} \cos\theta \right)$ $= \sqrt{7^2+24^2} \sin(\theta - A)$ $r = \sqrt{7^2+24^2}$ $= 25 \dots\dots\dots$ $A = \tan^{-1} \frac{24}{7}$ $\doteq 73.7^\circ \text{ (} 73^\circ 42' \text{ or } 1.29 \text{ rad.)}$	1A 1A 1A 1A	<u>Alternative Solutions:</u> $r\sin(\theta - A)$ $= r\sin\theta\cos A - r\cos\theta\sin A$ 1A $= 7\sin\theta - 24\cos\theta$ $r\cos A = 7$ ) ..... 1A $r\sin A = 24$ ) ..... $r = 25$ 1A $A = 73.7^\circ$ ..... 1A
(ii)	$y = 2(7\sin\theta - 24\cos\theta) + 14$ $= 2[25\sin(\theta - 73.7^\circ)] + 14$ $-1 \leq \sin(\theta - 73.7^\circ) \leq 1$ $-36 \leq y \leq 64 \dots\dots\dots$ <p>When <math>y = 64</math>,</p> $\sin(\theta - 73.7^\circ) = 1$ $\theta - 73.7^\circ = 180n^\circ + (-1)^n 90^\circ \text{ or } 360n^\circ + 90^\circ$ $\theta = 180n^\circ + (-1)^n 90^\circ + 73.7^\circ \text{ or } 360n^\circ + 163.7^\circ$	2M 1M+1M 1A+1A  1A  <u>1A</u> <u>12</u>	<u>Alternative Solution:</u> $y' = 50\cos(\theta - 1.29) = 0$ 1M $y'' = -50\sin(\theta - 1.29)$ 1M Max. $y = 64$ ..... 1A Min. $y = -36$ ..... 1A
(b)	$\cos\alpha \cos\beta = \frac{1}{6} \dots\dots\dots$ $\cos\alpha + \cos\beta = \frac{5}{6}$ $\left(\cos \frac{\alpha+\beta}{2} + \cos \frac{\alpha-\beta}{2}\right)^2 = \left(2\cos \frac{\alpha}{2} \cos \frac{\beta}{2}\right)^2$ $= (2\cos^2 \frac{\alpha}{2})(2\cos^2 \frac{\beta}{2}) \dots\dots\dots$ $= (1 + \cos\alpha)(1 + \cos\beta)$ $= 1 + \cos\alpha \cos\beta + \cos\alpha + \cos\beta$ $= 1 + \frac{1}{6} + \frac{5}{6} \dots\dots\dots$ $= 2$ $\cos \frac{\alpha+\beta}{2} + \cos \frac{\alpha-\beta}{2} = \sqrt{2}$	1 1 2A 1A 1A 1A 1M  8	
<u>Alternative Solution:</u> $\left(\cos \frac{\alpha+\beta}{2} + \cos \frac{\alpha-\beta}{2}\right)^2$ $= \cos^2 \frac{\alpha+\beta}{2} + \cos^2 \frac{\alpha-\beta}{2} + 2\cos \frac{\alpha+\beta}{2} \cos \frac{\alpha-\beta}{2}$ $= \frac{1}{2}[1 + \cos(\alpha+\beta)] + \frac{1}{2}[1 + \cos(\alpha-\beta)] + \cos\alpha + \cos\beta$ $= 1 + \cos\alpha + \cos\beta + \frac{1}{2}[\cos(\alpha+\beta) + \cos(\alpha-\beta)]$ $= 1 + \cos\alpha + \cos\beta + \cos\alpha \cos\beta \dots\dots\dots$ $= 1 + \frac{1}{6} + \frac{5}{6}$ $= 2$ $\cos \frac{\alpha+\beta}{2} \cos \frac{\alpha-\beta}{2} = \sqrt{2}$		1A 1A+1A 2A 1M	