

12. (b)




Figure 4

Figure 4 shows a circular pool of radius r metres centred at O . Two men, X and Y , holding the ends of a long rod, are walking in the direction shown at a speed of u metres per second. At a certain instant, the portion BD of the rod subtends an angle of 2θ radians at O and is at a distance ℓ metres from the midpoint C of the rim BD of the pool.

(i) Express ℓ in terms of r and θ .

Let A square metres be the area of the shaded region.

(ii) Express A in terms of r and θ .

(iii) Let $\frac{dA}{dt}$ (in m^2s^{-1}) be the rate of change of the area of the shaded region with respect to time.

Express $\frac{dA}{dt}$ in terms of r , θ and u . (Hint: $\frac{d\ell}{dt} = u$.)

Hence deduce that $\frac{dA}{dt} = ru$ when $\theta = \frac{\pi}{6}$.

(10 marks)

END OF PAPER

88-CE
A MATHS
PAPER II

HONG KONG EXAMINATIONS AUTHORITY
HONG KONG CERTIFICATE OF EDUCATION EXAMINATION 1988

附加數學 試卷二
ADDITIONAL MATHEMATICS PAPER II

11.15 am–1.15 pm (2 hours)
This paper must be answered in English

Answer ALL questions in Section A and any THREE questions from Section B.

All working must be clearly shown.

Unless otherwise specified in a question, it is sufficient for numerical answers to be given correct to three significant figures.

SECTION A (39 marks)

Answer ALL questions in this section.

1. Given $(1 + 3x)^4 (1 - 2x)^5 = 1 + ax + bx^2 + \text{higher powers of } x$, find the values of the constants a and b . (5 marks)

2. A and B are the points $(1, 2)$ and $(7, 4)$ respectively. P is a point on the line segment AB such that $\frac{AP}{PB} = k$.

- (a) Write down the coordinates of P in terms of k .
 (b) Hence find the ratio in which the line $7x - 3y - 28 = 0$ divides the line segment AB . (5 marks)




Figure 1

In Figure 1, the shaded region is bounded by the two curves $y = x(x + 3)$ and $y = x(5 - x)$. Find the area of the shaded region. (5 marks)

4. O and A are the points $(0, 0)$ and $(6, 0)$ respectively. $P(x, y)$ is a variable point such that $PO + PA = 10$. Find the equation of the locus of P , giving the answer in the form $ax^2 + by^2 + cx + d = 0$. (5 marks)

5. Prove, by mathematical induction, that

$$1^2 + 3^2 + \dots + (2n - 1)^2 = \frac{n(2n - 1)(2n + 1)}{3}$$

for all positive integers n . (6 marks)

6. Evaluate $\int_0^2 \frac{x^2 dx}{\sqrt{9 - x^3}}$. (6 marks)

7. (a) Without using calculators, show that $\frac{\pi}{10}$ is a root of $\cos 3\theta = \sin 2\theta$.
 (b) Given that $\cos 3\theta = 4 \cos^3 \theta - 3 \cos \theta$ and $\sin 2\theta = 2 \sin \theta \cos \theta$, find the value of $\sin \frac{\pi}{10}$, expressing the answer in surd form. (7 marks)

SECTION B (60 marks)

Answer any THREE questions from this section.
Each question carries 20 marks.

8. (a) Using the substitution $u = \sin x$, evaluate $\int_0^{\frac{\pi}{2}} \cos^7 x \, dx$.
Leave the answer as a fraction. (5 marks)

- (b) Let $y = \sin x \cos^{2n-1} x$, where n is a positive integer.

Find $\frac{dy}{dx}$.

Hence show that

$$2n \int \cos^{2n} x \, dx - (2n-1) \int \cos^{2n-2} x \, dx = \sin x \cos^{2n-1} x + C,$$

where C is a constant.

(4 marks)

- (c) (i) Using (b), show that

$$\int_0^{\frac{\pi}{2}} \cos^{2n} x \, dx = \frac{2n-1}{2n} \int_0^{\frac{\pi}{2}} \cos^{2n-2} x \, dx,$$

where n is a positive integer.

- (ii) Evaluate $\int_0^{\frac{\pi}{2}} \cos^6 x \, dx$ in terms of π .

(7 marks)

- (d) Evaluate $\int_0^{\frac{\pi}{2}} \sin^6 x \, dx$ in terms of π .

(4 marks)

9.




Figure 2

A balloon B is observed simultaneously from two points P and Q on a horizontal ground, P being at a distance c metres due north of Q . The bearings of the balloon from P and Q are $S\alpha^\circ E$ and $N\beta^\circ E$ respectively. The angle of elevation of B from P is θ° . R is the projection of B on the ground (see Figure 2).

- (a) Show that the balloon is at a height h metres where

$$h = \frac{c \tan \theta^\circ \sin \beta^\circ}{\sin(\alpha^\circ + \beta^\circ)}.$$

(6 marks)

- (b) Given $\theta = 40^\circ$, $\alpha = 54^\circ$ and $\beta = 46^\circ$,

- (i) find the angle of elevation of B from Q ;

- (ii) find the angle of elevation and the bearing of B from M , where M is the mid-point of PQ .

(14 marks)

10. (a) Using the substitution $x = a \sin \theta$, evaluate $\int_0^a \sqrt{a^2 - x^2} dx$,

where $a > 0$.

Hence, or otherwise, find the area of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$.

(6 marks)

- (b) (i) A pebble is in the shape of the solid of revolution of the ellipse $x^2 + \frac{y^2}{\left(\frac{3}{4}\right)^2} = 1$ about the x -axis.

Find the volume of the pebble.

(ii)




Figure 3

Figure 3 shows the cross-section of a bowl generated by revolving the lower half of the ellipse $\frac{x^2}{(2b)^2} + \frac{y^2}{b^2} = 1$ about the y -axis. The bowl contains water to a depth of h ($h < b$).

- (1) Show that the volume V of water in the bowl is given by $V = \frac{4}{3} \pi h^2 (3b - h)$.

Find $\frac{dV}{dh}$ when $h = \frac{b}{2}$.

- (2) Now the pebble in (i) is dropped into the bowl and is completely immersed in water. If $b = 5$ units and $h = 2.5$ units, show that the rise of water level is approximately equal to 0.01 unit. (14 marks)

11.




Figure 4

In Figure 4, S is the centre of the circle C which passes through $H(-3, 6)$ and touches the line $x - 5y + 59 = 0$ at $K(1, 12)$.

- (a) Find the coordinates of S .

Hence, or otherwise, find the equation of the circle C .

(8 marks)

The line $L: 3x - 2y - 5 = 0$ cuts the circle C at A and B .

- (b) Write down the equation of the family of circles through A and B .

Hence find the equation of the circle with AB as diameter.

(7 marks)

- (c) Show that $\angle ASB = 90^\circ$.

If P is any point on the circle C other than A or B , write down the two possible values of $\angle APB$.

(5 marks)

12. L is a line through the point $P(-2, 0)$ with slope m ($m \neq 0$), meeting the parabola $y^2 = 8x$ at the points $A(x_1, y_1)$ and $B(x_2, y_2)$.

- (a) Show that x_1 and x_2 are the roots of the equation

$$m^2 x^2 + (4m^2 - 8)x + 4m^2 = 0. \quad (3 \text{ marks})$$

- (b) Find $(x_1 - x_2)^2$ in terms of m . (5 marks)

- (c) Show that $AB^2 = \frac{64(1+m^2)(1-m^2)}{m^4}$. (5 marks)

- (d) Find the values of m for which L touches $y^2 = 8x$. (3 marks)

- (e) If $m = \frac{\sqrt{3}}{3}$ and C is the point $(2, 0)$, find by using (c), the area of $\triangle ABC$. (Leave the answer in surd form.) (4 marks)

END OF PAPER

SECTION A

Paper I

Candidates' performance on individual questions:

- Q.1** This was a straightforward question but candidates' performance was surprisingly poor. Wrongly memorizing the formula for differentiating $\operatorname{cosec} x$, quite a number of candidates made mistakes as follows:

(i) $f'(x) = 2 \operatorname{cosec} 3x \operatorname{cosec} 3x \cot 3x \cdot 3,$

(ii) $f'(x) = -2 \operatorname{cosec} 3x \cot^2 3x,$

(iii) $f'(x) = -2 \operatorname{cosec} 3x \operatorname{cosec} 3x \cot 3x.$

However, those who got the correct derivative of $f(x)$ could score full marks.

- Q.2** Most candidates performed well in the first part of this question. But in finding the second derivative, a great number of candidates made the following mistakes:

(i) $\frac{d^2 y}{dx^2} = \frac{-(-\sin y)}{(1 + \cos y)^2}$ in which $\left(\frac{dy}{dx}\right)$ was missing,

(ii) $\frac{d^2 y}{dx^2} = \frac{-\sin y \frac{dy}{dx}}{(1 + \cos y)^2}$ in which the minus sign was missing.

- Q.3** Candidates' performance was far from satisfactory. Many of them could not observe the relation between the two parts in this question. They solved the second part from the very beginning by putting $z_1 = x_1 + iy_1$ and $z_2 = x_2 + iy_2$ and were thus unable to go any further.

- Q.4** This question was well answered. Most candidates were able to score high marks. Others failed to do so mainly because mistakes were made in manipulation.

- Q.5** The candidates, in general, performed fairly well. A few of them overlooked the fact that the roots of the given equation

Additional Mathematics I

1. (a) Δx
 (b) $\frac{1}{2\sqrt{x+1}}$
2. $4x - 3y + 2 = 0$
 $2x - 3y - 5 = 0$
3. (a) $\frac{3}{13} + \frac{11}{13}i$
 (b) 74.7°
4. (a) $\frac{dy}{dx} = \cos x + 2 \sin x$
 $\frac{d^2y}{dx^2} = -\sin x + 2 \cos x$
 -2.24
5. $16m^2 - 16m - 60$
 $-\frac{3}{2} < m < \frac{5}{2}$
6. $a \cdot c = 6 + 4k$
 $b \cdot c = 16 + 6k$
 $k = 2$
7. $x < 0$ or $x > 1$
8. (a) (1.5, 1)
 (b) (i) x -intercept $= -2 \pm \sqrt{6}$
 y -intercept $= -0.5$
 (ii) $x > 4$ or $x < -1$
 (iii) (4, 1.5) is a maximum point.
 (-1, -1) is a minimum point.
 (d) (i) $y = 1 + \frac{4x-6}{x^2+4}$
9. (a) $\vec{AB} = -7i - 4j$
 $\vec{OC} = -i - 12j$
 (b) (i) $\vec{OX} = \frac{1}{1-k}(i + 2j)$
- (ii) $k = 1\frac{1}{2}$
 $\vec{AX} + \vec{BX} + \vec{CX} = 0$
 $\vec{AM} = -\frac{9}{2}i - 9j$
10. (a) (i) $PQ = 2\sqrt{2}$
 $RS = 2\sqrt{6}$
 (ii) x -coordinate of the mid-point of $RS = k$
 $k = -1$
 (b) $k = 3$ or -5
 $k = 3, (1, 2)$
 $k = -5, (-3, 2)$
 (c) $k > 3$ or $k < -5$
11. (a) (ii) $z = \cos\left(\frac{n\pi}{2} + \frac{\pi}{6}\right) + i \sin\left(\frac{n\pi}{2} + \frac{\pi}{6}\right)$
 $n = 0, 1, 2, 3$
 (b) (i) $\cos \frac{\pi}{3} + i \sin \frac{\pi}{3}$
 or $\cos\left(-\frac{\pi}{3}\right) + i \sin\left(-\frac{\pi}{3}\right)$
 (ii) (1) $x^2 - 2x + 1 = 0$
 (2) $x^2 + x + 1 = 0$
 (3) $x^2 + x + 1 = 0$
12. (a) (i) $\frac{\phi}{2\pi}$
 (iii) $\phi = \frac{2\sqrt{6}}{3}\pi$
 (b) (i) $r - r \cos \theta$
 (ii) $r^2 \theta - \frac{1}{2}r^2 \sin 2\theta$
 (iii) $2ru \sin \theta$

Additional Mathematics II

1. $a = 2$
 $b = -26$
2. (a) $\left(\frac{7k+1}{k+1}, \frac{4k+2}{k+1}\right)$
 (b) $3 : 1$
3. $\frac{1}{3}$
4. $16x^2 + 25y^2 - 96x - 256 = 0$
6. $\frac{4}{3}$
7. (b) $\frac{\sqrt{5}-1}{4}$
8. (a) $\frac{16}{35}$
 (b) $2n \cos^2 n x - (2n-1) \cos^2 n^{-2} x$
 (c) $\frac{5}{32}\pi$
 (d) $\frac{5}{32}\pi$
9. (b) (i) 36.7°
 (ii) $N83.2^\circ E$
10. (a) $\frac{\pi a^2}{4}$
 πab
 (b) (i) $\frac{3}{4}\pi$
 (ii) (1) $3\pi b^2$
11. (a) (2, 7)
 $x^2 + y^2 - 4x - 14y + 27 = 0$
 (b) $x^2 + y^2 - 4x - 14y + 27 + k(3x - 2y - 5) = 0$
 $x^2 + y^2 - 10x - 10y + 37 = 0$
 (c) $45^\circ, 135^\circ$
12. (b) $\frac{64(1-m^2)}{m^4}$
 (d) ± 1
 (e) $16\sqrt{2}$