

ADDITIONAL MATHEMATICS PAPER II

11.15 am-1.15 pm (2 hours)

This paper must be answered in English

Answer ALL questions in Section A and any
THREE questions in Section B.

All working must be clearly shown.

Unless otherwise specified in a question,
numerical answers must be given in exact value.

Section A (42 marks)

Answer ALL questions in this section.

1. Given that $(1 + x + ax^2)^8 = 1 + 8x + k_1x^2 + k_2x^3 + \dots$ terms involving higher powers of x .

(a) Express k_1 and k_2 in terms of a .

(b) If $k_1 = 4$, find the value of a .

Hence find the value of k_2 .

(5 marks)

2. Evaluate $\int_0^{\frac{\pi}{2}} (\sin x + \cos x)^2 dx$.

(5 marks)

3. Find the general solution of

$$\cos 4\theta + \cos 2\theta = \cos \theta.$$

(5 marks)

4. A family of straight lines is given by the equation

$$(2 - k)x + (1 + 2k)y - (4 + 3k) = 0,$$

where k is any constant.

Find equations of the two lines in the family whose distances from the origin are equal to 1.

(5 marks)

5. The slope at any point (x, y) of a curve C is given by

$$\frac{dy}{dx} = 4 - 2x$$

and C passes through the point $(1, 0)$.

- (a) Find an equation of C .
 (b) Find the area of the finite region bounded by C and the x -axis. (7 marks)

6.




Figure 1

In Figure 1, $PABCD$ is a right pyramid with a square base of sides of length 4 cm. $\angle PAB = 60^\circ$. Find, correct to the nearest 0.1 degree,

- (a) the angle between the plane PAB and the base $ABCD$,
 (b) the angle between the planes PAB and PAD . (7 marks)
7. (a) Prove, by mathematical induction, that

$$1^2 + 2^2 + \dots + n^2 = \frac{1}{6}n(n+1)(2n+1)$$

for all positive integers n .

- (b) Using the formula in (a), find the sum

$$1 \times 2 + 2 \times 3 + \dots + n(n+1). \quad (8 \text{ marks})$$

Section B (48 marks)

Answer any **THREE** questions in this section.

Each question carries 16 marks.

8. (a) Given that $\tan x = k \tan y$.
 (i) Show that $\sin(x + y) = (k + 1) \cos x \sin y$.
 (ii) Hence show that

$$(k + 1) \sin(x - y) = (k - 1) \sin(x + y). \quad (6 \text{ marks})$$

- (b) Given that

$$\tan(\theta + 10^\circ) = k \tan(\theta - 20^\circ)$$

has solutions in θ .

- (i) Show that $\sin(2\theta - 10^\circ) = \frac{k+1}{2(k-1)}$.
 (ii) Hence find the range of possible values of k . (7 marks)
- (c) Find the general solution of

$$\tan(\theta + 10^\circ) = -2 \tan(\theta - 20^\circ),$$

giving the answer correct to the nearest 0.1 degree.

(3 marks)

9. L is a straight line which passes through point $A(1, 2)$ and makes an angle θ with the positive x -axis. $P(x, y)$ is a point on L such that $AP = s$, as shown in Figure 2 (a).

- (a) Write down the coordinates of P in terms of s and θ . (2 marks)




Figure 2(a)

The circle $C : x^2 + y^2 - 6x - 10y + 30 = 0$ cuts the line L at points H and K (see Figure 2 (b)). Let $AH = s_1$, $AK = s_2$.

- (b) Show that s_1 and s_2 are the roots of the equation

$$s^2 - (4 \cos \theta + 6 \sin \theta) s + 9 = 0.$$

(3 marks)




Figure 2(b)

- (c) Using the result of (b), show that

$$HK^2 = 48 \sin \theta \cos \theta - 20 \cos^2 \theta. \quad (4 \text{ marks})$$

- (d) Using the result of (c), find equations of the two tangents from the point A to the circle C . (7 marks)

10. Given a hyperbola $H : \frac{x^2}{16} - \frac{y^2}{9} = 1$ and a straight line $L : 5x - 4y = 0$.

- (a) A and B are two points on H . The tangents to H at A and B are parallel to L . Find the coordinates of A and B . (7 marks)

- (b) Let $y = \frac{5}{4}x + c$ be an equation of a straight line which cuts H at two points P and Q . Find the coordinates of the mid-point of PQ in terms of c . (4 marks)

- (c) A chord of H is parallel to L and M is its mid-point. Using the result of (b), find an equation of the locus of M . Sketch the locus. (5 marks)

11.




Figure 3(a)

An object S is in the shape of the solid of revolution of the region bounded by the curve $x^2 = 4y$ and the line $y = 4$ revolved about the y -axis, as shown in Figure 3 (a).

- (a) Find the volume of S . (3 marks)

- (b) It is given that if S is cut by a plane parallel to its top surface at a distance h from O (see Figure 3 (a)), the mass of the part of S below the plane is given by

$$\int_0^h \pi(16y - 3y^2) dy, \text{ where } 0 < h \leq 4.$$

- (i) Find the mass of S .
- (ii) If the plane mentioned above cuts S into two parts of equal volumes, find
- (1) the value of h ,
 - (2) the ratio of the mass of the lower part to that of the upper part.
- (9 marks)




Figure 3(b)

In Figure 3(b), the shaded region is bounded by the curves $x^2 = 4y$, $x^2 = 4(y + t)$ and the line $y = 4$, where t is a small positive number. The solid of revolution of the shaded area revolved about the y -axis represents a layer of paint coated on the curved surface of S . Show that the volume of paint is approximately equal to $16\pi t$.

(4 marks)

12. Let m, n be positive integers.

(a) Given that $y = (1 + x)^{m+1} (1 - x)^n$. Find $\frac{dy}{dx}$.

Hence show that

$$(m + 1) \int (1 + x)^m (1 - x)^n dx = (1 + x)^{m+1} (1 - x)^n + n \int (1 + x)^{m+1} (1 - x)^{n-1} dx. \quad (3 \text{ marks})$$

- (b) Using the result of (a), show that

$$\int_{-1}^1 (1 + x)^m (1 - x)^n dx = \frac{n}{m+1} \int_{-1}^1 (1 + x)^{m+1} (1 - x)^{n-1} dx. \quad (3 \text{ marks})$$

- (c) Without using a binomial expansion, evaluate

$$\int_{-1}^1 (1 + x)^8 dx. \quad (2 \text{ marks})$$

- (d) Using the substitution $x = \tan \theta$, show that

$$\int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{\cos^2 2\theta (1 + \tan \theta)^4}{\cos^6 \theta} d\theta = \int_{-1}^1 (1 + x)^6 (1 - x)^2 dx.$$

Hence, using the results of (b) and (c), evaluate

$$\int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{\cos^2 2\theta (1 + \tan \theta)^4}{\cos^6 \theta} d\theta. \quad (8 \text{ marks})$$

END OF PAPER

1991

Additional Mathematics I

1. $2 + 3i$
2. $-\frac{1}{(1+x)^2}$
3. $-3, -1, 2$
4. (a) $1 + 2\cos 2x, -4\sin 2x$
 (b) $\frac{\pi}{3} + \frac{\sqrt{3}}{2}, \frac{2\pi}{3} - \frac{\sqrt{3}}{2}$
5. (a) $\frac{a+3b}{4}$
 (b) (i) $\frac{k+1}{4k}(a+3b)$
 (ii) $\frac{a+2mb}{1+m}$
 $\frac{5}{3} \cdot \frac{3}{2}$
6. (a) $y = 2$
 $x = 1$
7. (a) $1 - 2k + k^2$
 (b) $x^2 - (2-k)x + (1-2k+k^2) = 0$
 $0 \leq k \leq \frac{4}{3}$
8. (a) $(3-x)i - (y+1)j$
 $(x-7)i + (y-1)j$
 $(x-10)i + yj$
 (b) (ii) (1) 4, 2
9. (a) $-2(x+3)^2 - 5$
 (b) (i) $1, -\frac{3}{10}$
 (c) $\frac{3}{10}, -1$
10. (a) (i) $\cos \theta + i \sin \theta$, where $\theta = \pm \frac{2\pi}{3}$
 (ii) $\cos \theta + i \sin \theta$, where $\theta = \pm \frac{2\pi}{9}, \pm \frac{4\pi}{9}, \pm \frac{8\pi}{9}$
11. (c) (i) $h > 9$
 (ii) $18\sqrt{3}$
 (d) (ii) $p > 18\sqrt{3}$
12. (a) (i) $2\cos \phi$
 (b) (i) $\frac{\sin \theta}{2 \sin \phi}$
 (ii) $4 \sin \theta \sqrt{1 - \frac{1}{4} \cos^2 \theta}$
 (c) $\frac{-\sqrt{5}}{20} s^{-1}$

1991

Additional Mathematics II

1. (a) $8a + 28, 56a + 56$
 (b) $-3, -112$
2. $\frac{\pi}{2} + 1$
3. $2n\pi \pm \frac{\pi}{2}, \frac{2n\pi}{3} \pm \frac{\pi}{9}$
4. $x = 1$
 $3x - 4y + 5 = 0$
5. (a) $y = -x^2 + 4x - 3$
 (b) $1\frac{1}{3}$
6. (a) 54.7°
 (b) 109.5°
7. (b) $\frac{1}{3}n(n+1)(n+2)$
8. (b) (ii) $k \geq 3$ or $k \leq \frac{1}{3}$
 (c) $90n^\circ + 5^\circ + (-1)^n 4.8^\circ$
9. (a) $(1 + r \cos \theta, 2 + r \sin \theta)$
 (d) $x = 1$
 $5x - 12y + 19 = 0$
10. (a) $(5, \frac{9}{4}), (-5, -\frac{9}{4})$
 (b) $(-\frac{5c}{4}, \frac{-9c}{16})$
 (c) $y = \frac{9x}{20}$ ($x > 5$ or $x < -5$)
11. (a) 32π
 (b) (i) 64π
 (ii) (1) $2\sqrt{2}$
 (2) $(2\sqrt{2} - 1) : 1$
12. (a) $(m+1)(1+x)^m(1-x)^n - n(1+x)^{m+1}(1-x)^{n-1}$
 (c) $\frac{512}{9}$
 (d) $\frac{128}{63}$