



Solution	Marks	Remarks
1. $\int \cos^2 \theta d\theta$ $= \int \frac{1}{2}(1 - \cos 2\theta) d\theta$ $= \frac{1}{2}\theta + \frac{1}{4}\sin 2\theta + c$ (c is a constant)	1A 1M+1A <u>3</u>	1M for $\int \cos k\theta d\theta = \frac{1}{k} \sin k\theta$ withhold 1A if c was omitted
2. $\frac{(x+\Delta x)^3 - x^3}{\Delta x} = \frac{x^3 + 3x^2(\Delta x) + 3x(\Delta x)^2 + (\Delta x)^3 - x^3}{\Delta x}$ $= 3x^2 + 3x(\Delta x) + (\Delta x)^2$	1A 1A	For expanding $(x+\Delta x)^3$
$\frac{(x+\Delta x)^3 - x^3}{\Delta x} = \frac{[(x+\Delta x) - x][(x+\Delta x)^2 + (x+\Delta x)x + x^2]}{\Delta x}$ $= (x+\Delta x)^2 + (x+\Delta x)x + x^2$	1A 1A	
$\frac{d}{dx} x^3 = \lim_{\Delta x \rightarrow 0} \frac{(x+\Delta x)^3 - x^3}{\Delta x}$ $= \lim_{\Delta x \rightarrow 0} [3x^2 + 3x(\Delta x) + (\Delta x)^2]$ $= 3x^2$	1A <u>1A</u> <u>4</u>	} no mark if $\lim_{\Delta x \rightarrow 0}$ was omitted


Solution	Marks	Remarks
3. $\begin{cases} \alpha + \beta = 5 \\ \alpha\beta = k \end{cases}$ $ \alpha - \beta = 3$ $(\alpha + \beta)^2 = 3^2$ $\alpha^2 - 2\alpha\beta + \beta^2 = 9$ $(\alpha + \beta)^2 - 4\alpha\beta = 9$ $5^2 - 4k = 9$ $k = 4$	1M 1M 1M 1A	
<u>Alternative Solution (1)</u> $x = \frac{5 \pm \sqrt{25 - 4k}}{2}$ $ \alpha - \beta = 3$ $\left \frac{5 + \sqrt{25 - 4k}}{2} - \frac{5 - \sqrt{25 - 4k}}{2} \right = 3$ $ \sqrt{25 - 4k} = 3$ $25 - 4k = 3^2$ $k = 4$	1M 1M 1M 1A	Accept without absolute sign
<u>Alternative Solution (2)</u> $\begin{cases} \alpha + \beta = 5 \\ \alpha\beta = k \end{cases}$ $ \alpha - \beta = 3$ $\alpha - \beta = 3 \quad \text{or} \quad \alpha - \beta = -3$ $\begin{cases} \alpha + \beta = 5 \\ \alpha - \beta = 3 \end{cases} \quad \text{OR} \quad \begin{cases} \alpha + \beta = 5 \\ \alpha - \beta = -3 \end{cases}$ $\alpha = 4 \quad \text{and} \quad \beta = 1 \quad \alpha = 1 \quad \text{and} \quad \beta = 4$ $k = \alpha\beta$ $= 4$	1M 1M 1M 1A	
	<hr/> 4	

Solution	Marks	Remarks
4. $3x^2 + 3y^2 - 2xy = 12$ $6x + 6y \frac{dy}{dx} - 2y - 2x \frac{dy}{dx} = 0$ $\frac{dy}{dx} = \frac{y-3x}{3y-x}$ Put $x = 2, y = 0$: $\frac{dy}{dx} = \frac{-6}{-2}$ $\frac{dy}{dx} = 3$	$1A+1A+1A$ 1A <hr/> 4	$1A$ for $\frac{d}{dx}(3x^2 + 3y^2)$, $1A$ for $\frac{d}{dx}(2xy)$, $1A$ for $\frac{d}{dx}(12)$

Solution	Marks	Remarks
5. $x^2 > x $ $ x ^2 > x $ $ x (x -1) > 0$ $x \neq 0$ and $ x -1 > 0$ $ x > 1$ $x > 1$ or $x < -1$	1M 1A 1A 1A	$ x > 1$ or $ x < 0$
<u>Alternative Solution (1)</u> $ x < x^2$ $-x^2 < x < x^2$ $x^2 + x > 0$ and $x^2 - x > 0$ $x(x+1) > 0$ and $x(x-1) > 0$ $x > 0$ or $x < -1$ and $x > 1$ or $x < 0$ Combining the two cases, $x > 1$ or $x < -1$.	1M 1A+1A 1A	
<u>Alternative Solution (2)</u> Consider the two cases: (1) $x > 0$, (2) $x < 0$. Case 1 ($x > 0$): The inequality becomes $x^2 > x$ $x(x-1) > 0$ Since $x > 0$, so $x > 1$. Case 2 ($x < 0$): The inequality becomes $x^2 > -x$ $x(x+1) > 0$ Since $x < 0$, so $x < -1$. Combining the two cases, $x > 1$ or $x < -1$.	1M 1A 1A 1A	Accept including $x = 0$.
<u>Alternative Solution (3)</u> $x^2 > x $ $x^4 > x^2$ $x^2(x^2-1) > 0$ $x \neq 0$ and $x^2-1 > 0$ $x^2 > 1$ $x > 1$ or $x < -1$	1M 1A 1A 1A	$x^2 > 1$ or $x^2 < 0$
<u>Alternative Solution (4)</u>  From the above graph, $x^2 > x $ when $x > 1$ or $x < -1$.	1M+1A 1A 1A	1M for sketching the two graphs For intersecting at $x = 1$ and $x = -1$
	4	

Solution	Marks	Remarks
<p>6. </p> <p>(a) $\overrightarrow{OP} = \frac{\overrightarrow{a} + 3\overrightarrow{b}}{4}$</p> <p>(b) $\overrightarrow{OC} = \frac{4}{3}\overrightarrow{OP}$ $= \frac{4}{3}\left(\frac{\overrightarrow{a} + 3\overrightarrow{b}}{4}\right)$ $= \frac{1}{3}\overrightarrow{a} + \overrightarrow{b}$ $\overrightarrow{BC} = \overrightarrow{OC} - \overrightarrow{OB}$ $= \frac{1}{3}\overrightarrow{a} + \overrightarrow{b} - \overrightarrow{b}$ $= \frac{1}{3}\overrightarrow{a}$ <div style="border: 1px dashed black; padding: 2px; display: inline-block;">$= \frac{1}{3}\overrightarrow{OA}$</div></p> <p>$\therefore OA$ is parallel to BC.</p>	<p>1A</p> <p>1M</p> <p>1A</p> <p>1M</p> <p>1</p> <hr/> <p>5</p>	<p>Omitting vector sign in most cases (pp-1)</p>
<p>7. For $n=1$, LHS = $\frac{1}{2}$</p> <p>RHS = $2 - \frac{1+2}{2^1} = \frac{1}{2} = \text{LHS}$</p> <p>$\therefore$ the statement is true for $n=1$.</p> <p>Assume $\frac{1}{2} + \frac{2}{2^2} + \frac{3}{2^3} + \dots + \frac{k}{2^k} = 2 - \frac{k+2}{2^k}$,</p> <p>where k is a positive integer.</p> <p>$\frac{1}{2} + \frac{2}{2^2} + \frac{3}{2^3} + \dots + \frac{k}{2^k} + \frac{k+1}{2^{k+1}}$ $= 2 - \frac{k+2}{2^k} + \frac{k+1}{2^{k+1}}$ $= 2 - \frac{2(k+2) - (k+1)}{2^{k+1}}$ $= 2 - \frac{(k+1)+2}{2^{k+1}}$</p> <div style="border: 1px dashed black; padding: 5px; display: inline-block;"> <p>The statement is also true for $n = k + 1$ if it is true for $n = k$. By the principle of mathematical induction, the statement is true for all positive integers n.</p> </div>	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <hr/> <p>5</p>	<p>Not awarded if any one of the above marks was withheld</p>

Solution	Marks	Remarks
8. (a) The equation of the family of straight line is $2x - 3y + 4 + \lambda(x + y - 3) = 0$, where λ is a constant, and $x - y - 3 = 0$	1A	OR $(x + y - 3) + \lambda(2x - 3y + 4) = 0$.
OR $h(2x - 3y + 4) + k(x - y - 3) = 0$, where h, k are constants	1A	
<u>Alternative Solution</u> $\begin{cases} 2x - 3y + 4 = 0 \\ x + y - 3 = 0 \end{cases}$ Solve the equations, $x = 1$ and $y = 2$. The point of intersection of L_1 and L_2 is $(1, 2)$. The equation of the family of straight line is $y - 2 = m(x - 1)$	1A	
(b) $2x - 3y + 4 + \lambda(x + y - 3) = 0$ $(2 + \lambda)x + (\lambda - 3)y + (4 - 3\lambda) = 0$ $\frac{ 4 - 3\lambda }{\sqrt{(2 + \lambda)^2 + (\lambda - 3)^2}} = 1$ $(4 - 3\lambda)^2 = (2 + \lambda)^2 + (\lambda - 3)^2$ $7\lambda^2 - 22\lambda - 3 = 0$ $\lambda = 3$ or $\frac{1}{7}$ Put $\lambda = 3$ and $\frac{1}{7}$ into (1), the equation of the two lines are $x - 1 = 0$ and $3x - 4y + 5 = 0$.	1M+1M	1M for distance formula, 1M for substituting $(0, 0)$ & distance = 1 Accept omitting absolute sign
<u>Alternative Solution</u> $mx - y + (2 - m) = 0$ $\frac{ 2 - m }{\sqrt{m^2 + 1}} = 1$ $(2 - m)^2 = m^2 + 1$ $m = \frac{3}{4}$ \therefore the equation of the line is $y - 2 = \frac{3}{4}(x - 1)$ i.e. $3x - 4y + 5 = 0$. Another line which is of distance 1 from the origin is $x - 1 = 0$.	1M+1M 1A 1A	Same as above
	<u>5</u>	


Solution	Marks	Remarks
<p>9.</p>  $\text{Area} = \int_0^4 (x_2 - x_1) dy$ $= \int_0^4 \left(\frac{y+4}{4} - y^{\frac{1}{2}} \right) dy$ $= \left[\frac{y^2}{8} + y - \frac{2}{3} y^{\frac{3}{2}} \right]_0^4$ $= \frac{4^2}{8} + 4 - \frac{2}{3} (4^{\frac{3}{2}}) - 0$ $= \frac{2}{3}$	<p>1M+1A+1A</p> <p>1M</p> <p>1A</p>	<p>1M for $A = \int_a^b x dy$,</p> <p>1A for integrand, 1A for limit</p> <p>For evaluating $\int \frac{y+4}{4} dy$ & $\int y^{\frac{1}{2}} dy$</p>
<p><u>Alternative Solution</u></p> $\text{Area} = \int_0^2 y_2 dx - \int_1^2 y_1 dx$ $= \int_0^2 x^2 dx - \int_1^2 (4x-4) dx$ <div style="border: 1px solid black; padding: 5px; margin: 5px 0;"> $\text{OR} = \int_0^2 x^2 dx - \frac{1}{2} (4)(2-1)$ </div> $= \left[\frac{x^3}{3} \right]_0^2 - [2x^2 - 4x]_1^2$ $= \frac{8}{3} - 2$ $= \frac{2}{3}$	<p>1M+1A+1A</p> <p>1M</p> <p>1A</p>	<p>1M for $A = \int_a^b y dx$,</p> <p>1A for any correct expression</p> <p>For evaluating all primitive function(s)</p>
<p><u>5</u></p>		



	Solution	Marks	Remarks
10. (a)	$\frac{\sin \alpha + \sin \beta}{\cos \alpha + \cos \beta} = \frac{2 \sin \frac{\alpha + \beta}{2} \cos \frac{\alpha - \beta}{2}}{2 \cos \frac{\alpha + \beta}{2} \cos \frac{\alpha - \beta}{2}}$ $= \tan \frac{\alpha + \beta}{2}$	1A 1	
(b)	$3 \sin \alpha - 4 \cos \alpha = 4 \cos \beta - 3 \sin \beta$ $3 \sin \alpha + 3 \sin \beta = 4 \cos \alpha + 4 \cos \beta$ $\frac{\sin \alpha + \sin \beta}{\cos \alpha + \cos \beta} = \frac{4}{3}$		
	Using (a), $\tan \left(\frac{\alpha + \beta}{2}\right) = \frac{4}{3}$	1M	
	$\tan (\alpha + \beta) = \frac{2 \tan \left(\frac{\alpha + \beta}{2}\right)}{1 - \tan^2 \left(\frac{\alpha + \beta}{2}\right)}$		
	$= \frac{2 \left(\frac{4}{3}\right)}{1 - \left(\frac{4}{3}\right)^2}$	1M	
	$= -\frac{24}{7}$	1A	
		5	

Solution	Marks	Remarks
11. $\left(\frac{1+3i}{1-2i}\right)^{20}$ $\frac{1+3i}{1-2i} = \frac{1+3i}{1-2i} \left(\frac{1+2i}{1+2i}\right)$ $= \frac{1-2i+3i-6}{1+4}$ $= -1+i$ $= \sqrt{2}\left(\cos\frac{3\pi}{4} + i\sin\frac{3\pi}{4}\right)$ $\left(\frac{1+3i}{1-2i}\right)^{20} = \left[\sqrt{2}\left(\cos\frac{3\pi}{4} + i\sin\frac{3\pi}{4}\right)\right]^{20}$ $= (\sqrt{2})^{20} \left[\cos\left(\frac{3\pi}{4} \times 20\right) + i\sin\left(\frac{3\pi}{4} \times 20\right)\right]$ $= (\sqrt{2})^{20} (\cos \pi + i \sin \pi)$ $= (\sqrt{2})^{20} (-1)$ $= -2^{10} \quad (= -1024)$	1M 1A 1M 1M 1A	For expressing in polar form Accept $2^{10}(\cos \pi + i \sin \pi)$
<u>Alternative Solution</u> $\frac{1+3i}{1-2i} = \dots = -1+i$ $(-1+i)^2 = (-1)^2 - 2i + i^2$ $= -2i$ $\left(\frac{1+3i}{1-2i}\right)^{20} = (-2i)^{10}$ $= (-2)^{10} (-1)^5$ $= -2^{10}$	1M+1A 1M 1M 1A	same as above
	<hr/> 5	

Solution	Marks	Remarks
<p>12. $(2x^2 + \frac{1}{x})^9$</p> <p>General term $= {}_9C_r (2x^2)^{9-r} (\frac{1}{x})^r$ $= {}_9C_r (2^{9-r}) (x^{18-3r})$</p> <p>(a) Put $18 - 3r = 0$: $r = 6$ \therefore constant term $= {}_9C_6 (2^{9-6})$ $= 84(8) = 672$</p> <p>(b) Put $18 - 3r = 2$: $r = \frac{16}{3}$ which is not an integer. \therefore there is no x^2 term.</p>	<p>1A</p> <p>1M</p> <p>1M</p> <p>1A</p> <p>1A+1M</p>	<p>${}_9C_r (2x^2)^r (\frac{1}{x})^{9-r}$</p> <p>${}_9C_r (2^r) (x^{3r-9})$</p>
<p><u>Alternative Solution (1)</u></p> $(2x^2 + \frac{1}{x})^9$ $= (2x^2)^9 + {}_9C_1 (2x^2)^8 (\frac{1}{x}) + {}_9C_2 (2x^2)^7 (\frac{1}{x})^2$ $+ {}_9C_3 (2x^2)^6 (\frac{1}{x})^3 + {}_9C_4 (2x^2)^5 (\frac{1}{x})^4 + {}_9C_5 (2x^2)^4 (\frac{1}{x})^5$ $+ {}_9C_6 (2x^2)^3 (\frac{1}{x})^6 + {}_9C_7 (2x^2)^2 (\frac{1}{x})^7 + {}_9C_8 (2x^2)^1 (\frac{1}{x})^8$ $+ (\frac{1}{x})^9$ <hr style="border-top: 1px dashed black;"/> $= 2^9 x^{18} + 9(2^8) x^{15} + 36(2^7) x^{12} + 84(2^6) x^9$ $+ 126(2^5) x^6 + 126(2^4) x^3 + 84(2^3) + 36(2^2) x^{-3}$ $+ 9(2) x^{-6} + x^{-9}$ $= 512x^{18} + 2304x^{15} + 4608x^{12} - 5376x^9$ $+ 4032x^6 + 2016x^3 + 672 + 144x^{-3} + 18x^{-6} + x^{-9}$ <hr style="border-top: 1px dashed black;"/> <p>Constant term $= {}_9C_6 (2^3)$ $= 84(8) = 672$</p> <p>The above expansion indicates that there is no x^2 term.</p>	<p>1M+1A</p> <p>1M</p> <p>1A</p> <p>1A-1M</p>	
<p><u>Alternative Solution (2)</u></p> $(2x^2 + \frac{1}{x})^9 = \frac{(2x^3 + 1)^9}{x^9}$ <p>Constant term $= \frac{{}_9C_3 (2x^3)^3}{x^9}$ $= {}_9C_3 (2^3)$ $= 84(8) = 672$</p> <p>The numerator does not contain an x^{11} term, so there is no x^2 term.</p>	<p>1M</p> <p>1M+1A</p> <p>1A</p> <p>1A+1M</p>	
	<hr style="width: 50px; margin: auto;"/> <p>6</p>	

Solution	Marks	Remarks								
<p>13. $f(x) = 2 \sin x - x$ $f'(x) = 2 \cos x - 1$ $f'(x) = 0$ $2 \cos x - 1 = 0$ $\cos x = \frac{1}{2}$ $x = \frac{\pi}{3}$</p> <p>$f''(x) = -2 \sin x$ $f''(\frac{\pi}{3}) = -\sqrt{3} < 0$</p> <p>$\therefore f(x)$ attains a maximum at $x = \frac{\pi}{3}$.</p>	<p>1M</p> <p>1A</p> <p>1M+1A</p>	<p>Withhold this mark for $x = 60^\circ$</p>								
<p><u>Alternative Solution for checking</u> $f'(x) = 2 \cos x - 1$</p> <table border="1" data-bbox="199 750 805 873"> <tr> <td>x</td> <td>$0 \leq x < \frac{\pi}{3}$</td> <td>$\frac{\pi}{3}$</td> <td>$\frac{\pi}{3} < x \leq \pi$</td> </tr> <tr> <td>$f'(x) = 0$</td> <td>+ve</td> <td>0</td> <td>-ve</td> </tr> </table> <p>Since $f'(x)$ changes from +ve to -ve at $x = \frac{\pi}{3}$, $f(x)$ attains a maximum at $x = \frac{\pi}{3}$.</p>	x	$0 \leq x < \frac{\pi}{3}$	$\frac{\pi}{3}$	$\frac{\pi}{3} < x \leq \pi$	$f'(x) = 0$	+ve	0	-ve	<p>1M+1A</p>	
x	$0 \leq x < \frac{\pi}{3}$	$\frac{\pi}{3}$	$\frac{\pi}{3} < x \leq \pi$							
$f'(x) = 0$	+ve	0	-ve							
<p>$f(\frac{\pi}{3}) = 2 \sin \frac{\pi}{3} - \frac{\pi}{3}$ $= \sqrt{3} - \frac{\pi}{3}$</p> <p>Since $f(x)$ is continuous and has only one turning point, the greatest value of $f(x)$ is $\sqrt{3} - \frac{\pi}{3}$.</p> <p>The least value of $f(x)$ occurs at one of the end-points. $f(0) = 0$ $f(\pi) = -\pi$ \therefore the least value of $f(x)$ is $-\pi$.</p>	<p>1A</p> <p>1M</p> <p><u>1A</u> <u>7</u></p>	<p>Not awarded if checking was incomplete (can be omitted)</p>								


Solution	Marks	Remarks
<p>14.</p> 		
<p>(a) $\vec{a} \cdot \vec{b} = \vec{a} \vec{b} \cos \angle AOB$ $= 3(1) \cos 120^\circ$ $= -\frac{3}{2}$</p>	<p>1M <u>1A</u> <u>2</u></p>	
<p>(b) $\vec{OC} = \frac{2\vec{a} + 3\vec{b}}{2+3} = \frac{2}{5}\vec{a} + \frac{3}{5}\vec{b}$ $\vec{OD} = k\vec{OC}$ $= \frac{2k}{5}\vec{a} + \frac{3k}{5}\vec{b}$ $\vec{AD} = \vec{OD} - \vec{OA}$ $= \frac{2k}{5}\vec{a} + \frac{3k}{5}\vec{b} - \vec{a}$ $= (\frac{2k}{5} - 1)\vec{a} + \frac{3k}{5}\vec{b}$ $\vec{AD} \cdot \vec{AB} = 0$ $[(\frac{2k}{5} - 1)\vec{a} + \frac{3k}{5}\vec{b}] \cdot (\vec{b} - \vec{a}) = 0$ $(\frac{2k}{5} - 1)\vec{a} \cdot \vec{b} - (\frac{2k}{5} - 1)\vec{a} \cdot \vec{a} + \frac{3k}{5}\vec{b} \cdot \vec{b} - \frac{3k}{5}\vec{b} \cdot \vec{a} = 0$ $(\frac{2k}{5} - 1)(-\frac{3}{2}) - (\frac{2k}{5} - 1)(3)^2 + \frac{3k}{5}(1)^2 - \frac{3k}{5}(-\frac{3}{2}) = 0$ $\frac{105}{2} - \frac{27}{2}k = 0$ $k = \frac{35}{9}$</p>	<p>1M 1 1M 1M 1M <u>1A</u> <u>6</u></p>	<p>For distributive law For $\vec{a} \cdot \vec{a} = 9$ or $\vec{b} \cdot \vec{b} = 1$</p>
<p>(c) $\vec{OC} \cdot \vec{OB} = (\frac{2}{5}\vec{a} + \frac{3}{5}\vec{b}) \cdot \vec{b}$ $= \frac{2}{5}\vec{a} \cdot \vec{b} + \frac{3}{5}\vec{b} \cdot \vec{b}$ $= \frac{2}{5}(-\frac{3}{2}) + \frac{3}{5}(1)^2 = 0$ $\therefore \angle BOC = \frac{\pi}{2}$ (OR $\vec{OC} \perp \vec{OB}$) $\angle BOC = \angle DAC = \frac{\pi}{2}$ $\angle BCO = \angle DCA$ (vertically opposite \angles) $\angle OBC = \angle ADC$ (\angle sum of Δ) $\therefore \triangle OCB \sim \triangle ACD$ (AAA)</p>	<p>1M 1A 1A 1</p>	<p>Omitting vector sign or dot product sign in most cases (pp-1)</p>


Solution	Marks	Remarks
<p>Alternative solution (1)</p>  $AB^2 = 3^2 + 1^2 - 2(3)(1)\cos 120^\circ$ $AB = \sqrt{13}$ $\cos \angle OAB = \frac{3^2 + (\sqrt{13})^2 - 1^2}{2(3)(\sqrt{13})} = \frac{7}{2\sqrt{13}}$ $AC = \frac{3}{5}(AB) = \frac{3\sqrt{13}}{5}$ $OC^2 = 3^2 - \left(\frac{3\sqrt{13}}{5}\right)^2 - 2(3)\left(\frac{3\sqrt{13}}{5}\right)\left(\frac{7}{2\sqrt{13}}\right) = \frac{27}{25}$ $OB^2 + OC^2 = 1^2 + \frac{27}{25}$ $= \frac{52}{25} = \left(\frac{2\sqrt{13}}{5}\right)^2 = BC^2$ <p>$\therefore OB \perp OC$ (OR $\angle BOC = \frac{\pi}{2}$)</p> <p>∴</p> <p>∴</p>	<p>1M</p> <p>1A</p> <p>1A+1</p>	<p>(same as above)</p>
<p>Alternative solution (2)</p>  $BC = \frac{2\sqrt{13}}{5}, AC = \frac{3\sqrt{13}}{5}$ $OC = \frac{3\sqrt{13}}{5}$ $CD = \left(\frac{35}{9} - 1\right) OC = \frac{26\sqrt{13}}{15}$ $\frac{OC}{BC} = \frac{3\sqrt{13}/5}{2\sqrt{13}/5} = \frac{3\sqrt{13}}{2\sqrt{13}}$ $\frac{AC}{CD} = \frac{3\sqrt{13}/5}{26\sqrt{13}/15} = \frac{3\sqrt{13}}{2\sqrt{13}} = \frac{OC}{BC}$ <p>$\angle BCO = \angle DCA$ (vertically opposite \angles)</p> <p>$\triangle OCB \sim \triangle ACD$ (ratio of 2 sides, incl. \angle)</p>	<p>1M+1A</p> <p>1A</p> <p>1</p>	
	<p>4</p>	

Solution	Marks	Remarks
15. (a) (i) $x^2 = 4y$ $2x = 4 \frac{dy}{dx}$ At S , $\frac{dy}{dx} = s$ Equation of L_1 $\frac{y - s^2}{x - 2s} = s$ $sx - y = s^2$	1M 1A	
<u>Alternative solution (1)</u> Using the formula $xx_1 = 2(y + y_1)$, equation of L_1 is $x(2s) = 2(y + s^2)$ $sx - y = s^2$	1A 1A	
<u>Alternative solution (2)</u> Let the equation of L_1 be $y - s^2 = m(x - 2s)$ $\begin{cases} x^2 = 4y \\ y - s^2 = m(x - 2s) \end{cases}$ $x^2 - 4mx + (8ms - 4s^2) = 0$ $\Delta = 16m^2 - 4(8ms - 4s^2) = 0$ $m = s$ \therefore the equation of L_1 is $y = sx - s^2$.	1M 1A	
(ii) Equation of L_2 is $tx - y = t^2$. $\begin{cases} sx - y = s^2 & \text{----- (1)} \\ tx - y = t^2 & \text{----- (2)} \end{cases}$ (1) - (2): $(s - t)x = s^2 - t^2$ $x = s + t$ Substitute $x = s + t$ into (1): $s(s + t) - y = s^2$ $y = st$ Since the two tangents meet at the point $P(\alpha, \beta)$, $\begin{cases} s + t = \alpha \\ st = \beta \end{cases}$	1M 1M 1M 1	For replacing s by t
<u>Alternative solution</u> Equation of L_2 is $tx - y = t^2$. Since $P(\alpha, \beta)$ lies on both tangents, s, t are the roots of the equation $z\alpha - \beta - z^2 = 0$ in z . $z^2 - \alpha z + \beta = 0$ $\therefore \begin{cases} s + t = \alpha \\ st = \beta \end{cases}$	1M 2M 1	

Solution	Marks	Remarks
<p>(iii) Slope of $L_1 = s$ Slope of $L_2 = t$ If the line makes equal angles with L_1 and L_2.</p> $\frac{s-1}{1+s} = \frac{1-t}{1+t} \quad \text{OR} \quad \left \frac{1-s}{1+s} \right = \left \frac{1-t}{1+t} \right $ <p>$(s-1)(1+t) = (1+s)(1-t)$ $st = 1$</p>	<p>1M+1M</p> <hr/> <p>1</p> <hr/> <p>9</p>	<p>1M for $\tan \theta = \frac{m_2 - m_1}{1 + m_2 m_1}$, 1M for substitution</p>
<p>(b) Let the coordinates of R be (α, β) and the two points of tangency be $(2s, s^2)$ and $(2t, t^2)$.</p> <p>From (a) (ii), $\begin{cases} s+t = \alpha \\ st = \beta \end{cases}$.</p> <p>From (a) (iii), $st = 1$.</p> <p>$\therefore \beta = st = 1$</p> <p>Since R lies on the line $x - y + 4 = 0$,</p> $\alpha - 1 + 4 = 0$ $\alpha = -3$ <p>\therefore the coordinates of R are $(-3, 1)$.</p>	<p>1A</p> <p>1M</p> <hr/> <p>1A</p> <hr/> <p>3</p>	

Solution	Marks	Remarks
16. (a)	1A 1M 1A 1A 4	Position of A and B Position of C Quadrilateral OACB (a // -gram with OA = OB) Label theta Not labelling the axes (pp-1)
(b) (i) $z_4 = z_2 - z_1$ $= (\cos \theta - 1) + i \sin \theta$ $z_3 = z_1 + z_2$ $= (1 + \cos \theta) + i \sin \theta$ $\frac{z_4}{z_3} = \frac{(\cos \theta - 1) + i \sin \theta}{(1 + \cos \theta) + i \sin \theta} \cdot \frac{(1 + \cos \theta) - i \sin \theta}{(1 + \cos \theta) - i \sin \theta}$ $= \frac{(\cos \theta - 1)(1 - \cos \theta) - i \sin \theta(\cos \theta - 1) + i \sin \theta(1 + \cos \theta) - \sin^2 \theta}{(1 + \cos \theta)^2 + \sin^2 \theta}$ $= \frac{\cos^2 \theta - 1 + i[-\sin \theta \cos \theta + \sin \theta + \sin \theta + \sin \theta \cos \theta] + \sin^2 \theta}{1 + 2 \cos \theta + \cos^2 \theta + \sin^2 \theta}$ $= \frac{i 2 \sin \theta}{2(1 + \cos \theta)}$ $= \frac{i \sin \theta}{1 + \cos \theta}$	1A 1M 1A 1	
<u>Alternative solution (1)</u> $z_4 = (\cos \theta - 1) + i \sin \theta$ $z_3 = (1 + \cos \theta) + i \sin \theta$ $z_4(1 + \cos \theta) = [(\cos \theta - 1) + i \sin \theta](1 + \cos \theta)$ $= (\cos \theta - 1)(1 + \cos \theta) + i \sin \theta(1 + \cos \theta)$ $= -\sin^2 \theta + i(\sin \theta + \sin \theta \cos \theta)$ $z_3(i \sin \theta) = [(1 + \cos \theta) + i \sin \theta]i \sin \theta$ $= i(\sin \theta + \sin \theta \cos \theta) - \sin^2 \theta$ $\therefore z_4(1 + \cos \theta) = z_3(i \sin \theta)$ $\frac{z_4}{z_3} = \frac{i \sin \theta}{1 + \cos \theta}$	1A 1M 1A 1	

Solution	Marks	Remarks
<p>Alternative solution (2)</p> $z_4 = (\cos \theta - 1) + i \sin \theta, z_3 = (1 + \cos \theta) + i \sin \theta$ $\frac{z_4}{z_3} = \frac{-1 + \cos \theta + i \sin \theta}{1 + \cos \theta + i \sin \theta}$ $= \frac{-2 \sin^2 \frac{\theta}{2} + 2i \sin \frac{\theta}{2} \cos \frac{\theta}{2}}{2 \cos^2 \frac{\theta}{2} + 2i \sin \frac{\theta}{2} \cos \frac{\theta}{2}}$ $= \frac{2i \sin \frac{\theta}{2} (\cos \frac{\theta}{2} + i \sin \frac{\theta}{2})}{2 \cos \frac{\theta}{2} (\cos \frac{\theta}{2} + i \sin \frac{\theta}{2})}$ $= i \tan \frac{\theta}{2}$ $= i \frac{\sin \frac{\theta}{2} \left(\frac{\cos \frac{\theta}{2}}{\cos \frac{\theta}{2}} \right)}{\cos \frac{\theta}{2} \left(\frac{\cos \frac{\theta}{2}}{\cos \frac{\theta}{2}} \right)}$ $= \frac{i \sin \theta}{1 + \cos \theta}$	<p>1A</p> <p>1M</p> <p>1A</p> <p>1</p>	
<p>$\arg \left(\frac{z_4}{z_3} \right) = \frac{\pi}{2}$</p>	<p>1A</p>	
<p>(ii) Let ϕ be the angle between OC and AB.</p>		
		
<p>$\arg \left(\frac{z_4}{z_3} \right) = \frac{\pi}{2}$</p>		
<p>$\arg(z_4) - \arg(z_3) = \frac{\pi}{2}$</p>	<p>1M</p>	<p>For $\arg \left(\frac{z_4}{z_3} \right) = \arg(z_4) - \arg(z_3)$</p>
<p>$\arg(z_4) = \beta,$ $\arg(z_3) = \alpha$ (see Figure above)</p>	<p>1A</p>	
<p>$\arg(z_4) - \arg(z_3) = \beta - \alpha$ $= \phi$</p>		
<p>$\therefore \phi = \frac{\pi}{2},$ i.e. the diagonals of $OACB$ are perpendicular to each other.</p>	<p>1</p> <hr/> <p>8</p>	

Solution	Marks	Remarks
17. (a) The co-ordinates of P are (a, b) .	1A 1	
(b) (i) $g(x) = (x-b)^2 + a$ Substitute $Q(b, a)$ into $y = f(x)$: $a = -(b-a)^2 + b$ $(b-a)^2 = b-a \dots (1)$ OR $a^2 - 2ab + b^2 + a - b = 0$ $g(a) = (a-b)^2 + a$ $= b - a + a$ OR $= a^2 - 2ab + b^2 + a$ $= b$ $\therefore y = g(x)$ passes through P .	1A 1A 1M 1M 1	For using (1)
<u>Alternative Solution</u> Substitute $Q(b, a)$ into $y = f(x)$: $a = -(b-a)^2 + b$ $(b-a)^2 = b-a$ $b-a = 0$ or $b-a = 1$ Case 1: $(b-a) = 0$, i.e. $b = a$ $g(a) = (a-a)^2 + a$ $= a = b$ Case 2: $(b-a) = 1$ $= g(a) = (a-b)^2 + a$ $= 1 + a = b$ $g(a) = b$ in both cases. $\therefore y = g(x)$ passes through P .	1A 1M+1M 1	1M for considering $g(a)$, 1M for using $b-a = 0$ or $b-a = 1$
(ii) Since $y = f(x)$ touches the x -axis, y -coordinate of vertex = 0, i.e. $b = 0$. Substitute $b = 0$ into (1): $(0-a)^2 = 0-a$ $a = 0$ or -1 Case 1: $a = 0, b = 0$ $f(x) = -x^2$ and $g(x) = x^2$	1M 1M	$-x^2 + 2ax + (b-a^2) = 0$ $\Delta = 4a^2 + 4(b-a^2) = 0$ $b = 0$
	1A+1A	Axes not labelled (pp-1)


Solution

Marks

Remarks

Case 2: $a = -1, b = 0$



$f(x) = -(x-1)^2$ and $g(x) = x^2 - 1$




1A+1A


Withhold 1A if $(-1, 0)$ or $(0, -1)$ was not labelled

11

Solution	Marks	Remarks
<p>18. (a)</p>  <p>Let S be the point on PQ such that $OS \perp PQ$ and $RS \perp PQ$.</p> <p>Area of $\triangle OPQ = \frac{1}{2}(PQ)(OS)$</p> <p>Area of $\triangle RPQ = \frac{1}{2}(PQ)(RS)$</p> $\frac{\text{Area of } \triangle OPQ}{\text{Area of } \triangle RPQ} = \frac{\frac{1}{2}(PQ)(OS)}{\frac{1}{2}(PQ)(RS)}$ $= \frac{OS}{RS}$ $= \cos \theta$	<p>1M</p> <p>1A</p> <p>1A</p> <hr/> <p>$\frac{1}{4}$</p>	
<p>(b) (i)</p>  <p>Let M be the foot of perpendicular from C to AB and $\angle CME = \phi$.</p> $\frac{1}{2}(AB)(CM) = 12$ $\frac{1}{2}(6)(CM) = 12$ $CM = 4$ $\sin \phi = \frac{CE}{CM}$ $= \frac{2}{4} = \frac{1}{2}$ $\phi = \frac{\pi}{6}$ <p>From (a), area of $\triangle EAB = (\text{area of } \triangle CAB) \cos \phi$</p> $= 12 \cos \frac{\pi}{6}$ $= 6\sqrt{3} \text{ m}^2$ <p>\therefore the area of the shadow is $6\sqrt{3} \text{ m}^2$.</p>	<p>1M</p> <p>1M</p> <p>1M</p> <p>1A</p>	<p>For finding CM</p> <p>For finding ϕ, $\cos \phi$ or $ME = (\sqrt{12})$</p> $\text{OR } = \frac{AB \times ME}{2} \quad \text{OR } = 12\sqrt{1 - \sin^2 \phi}$ $= \frac{6 \times \sqrt{12}}{2} \quad = 12\sqrt{1 - \left(\frac{1}{2}\right)^2}$ $= 6\sqrt{3} \text{ m}^2 \quad = \sqrt{144 - 36}$ $= 6\sqrt{3}$

Solution	Marks	Remarks
<p>(ii)</p>  <p>Let α be the angle between the board and the ground, h be the altitude of the board from the vertex fastened to the top of the pole.</p> <p>From (a), area of shadow = $12\cos\alpha$</p> <p>In order for the area of the shadow to be the greatest, $\cos\alpha$ should be the greatest.</p> <p>$\sin\alpha$ should be the least.</p> <p>h should be the greatest.</p> <p>Since $6 > x > y$, the altitude from B to CA is the longest among the 3 altitudes. So vertex B should be fastened to the top of the pole.</p>	<p>1M</p> <p>1M</p> <p>1M</p> <p>1A</p>	<p><u>OR</u> $\frac{2}{h}$ (or α) should be the least</p>
<p><u>Alternative Solution</u></p> <p>If vertex B is fastened to the top of the pole,</p> $h = \frac{12(2)}{y} = \frac{24}{y}$ $\sin\alpha = \frac{2}{24/y} = \frac{y}{12}$ $\cos\alpha = \sqrt{1 - \left(\frac{y}{12}\right)^2}$ <p>Area of shadow = $12\sqrt{1 - \left(\frac{y}{12}\right)^2} = \sqrt{144 - y^2}$</p> <p>Similarly, if vertex A is fastened to the top of the pole, area of shadow = $\sqrt{144 - x^2}$</p> <p>Since $6 > x > y$,</p> $\sqrt{144 - 6^2} < \sqrt{144 - x^2} < \sqrt{144 - y^2}$ <p>area of the largest shadow = $\sqrt{144 - y^2}$.</p> <p>So vertex B should be fastened to the top of the pole.</p>	<p>1M</p> <p>1A</p> <p>1M</p> <p>1A</p>	
	<p>8</p>	

Solution	Marks	Remarks
<p>19. (a) $V = \int_0^h \pi x^2 dy$</p> $= \int_0^h \pi y dy$ $= \left[\frac{\pi y^2}{2} \right]_0^h$ $V = \frac{1}{2} \pi h^2$ <p>At $x = 2, y = 2^2 = 4.$</p> <p>Capacity of the tank $= \frac{1}{2} \pi (4)^2$</p> $= 8\pi$ <p>Time required to fill the tank</p> $= \frac{8\pi}{2\pi}$ $= 4 \text{ (minutes)}$	<p>1M+1A</p> <p>1A</p> <p>1M</p> <p><u>1</u></p> <p><u>5</u></p>	<p>1M for $V = \int_a^b \pi x^2 dy$</p>
<p>(b) $V = \frac{1}{2} \pi h^2$</p> <p>Differentiate with respect to t:</p> $\frac{dV}{dt} = \frac{1}{2} \pi (2h) \frac{dh}{dt}$ <p>Put $\frac{dV}{dt} = 2\pi$:</p> $2\pi = \pi h \frac{dh}{dt}$ $\frac{dh}{dt} = \frac{2}{h}$	<p>1M</p> <p>1M</p> <p><u>1</u></p> <p><u>3</u></p>	
<p>(c) From (b), $\frac{dh}{dt} = \frac{2}{h}.$</p> <p>The rate of change of h decreases as t increases. So Sketch C best describes the variation of h with $t.$</p>	<p>1M+1A</p>	<p>1M for a correct argument</p>
<p><u>Alternative solution (1)</u></p> <p>Sketch B is incorrect as $\frac{dh}{dt}$ is not a constant. * the x-section of the tank is non-uniform.</p> <p>As h increases, the surface area increases. Since $\frac{dV}{dt}$ is a constant, h will increase at a lower rate. So Sketch C best describes the variation of h with $t.$</p>	<p>1M</p> <p>1A</p>	

Solution	Marks	Remarks
<p>Alternative solution (2)</p> $\frac{dh}{dt} = \frac{2}{h}$ $\frac{d^2h}{dt^2} = \frac{-2}{h^2} \frac{dh}{dt}$ $= \frac{-4}{h^3} < 0$ <p>So Sketch C best describes the variation of h with t.</p>	<p>1M</p> <p>1A</p>	
<p>Alternative solution (3)</p> $\frac{dh}{dt} = \frac{2}{h}$ $\int h dh = \int 2 dt$ $\frac{h^2}{2} = 2t + c$ <p>At $t=0, h=0 \therefore c=0$</p> $h^2 = 4t$ <p>So Sketch C best describes the variation of h with t.</p>	<p>1M</p> <p>1A</p>	
<p>(d)</p>  <p style="text-align: center;">Sketch C</p>	<p style="text-align: center;"><u>2</u></p> <p>1M+1A</p> <p style="text-align: center;"><u>2</u></p>	<p>1M for a straight line with +ve slope for $h > 2$ Withhold 1 mark if not drawn on the sketch chosen in (c)</p>